

Radiative and rare kaon decays : an update ¹

Giancarlo D'Ambrosio ^a

^aINFN, Sezione di Napoli, Dip. di Scienze Fisiche, Univ. di Napoli, I-80125 Napoli, Italy.

Abstract

We review some new developments in the theoretical description of the processes $K \rightarrow \pi\gamma\gamma$, $K_L \rightarrow \gamma\ell^+\ell^-$, $K_L \rightarrow \pi^0 e^+e^-$ and $K_L \rightarrow \mu^+\mu^-$.

1 Introduction

Radiative non-leptonic kaon decays may play a crucial role in our understanding of fundamental questions: the validity of the Standard Model (SM) (fixing the values of the CKM parameters), the origin of CP violation and as a χ PT (Chiral Perturbation Theory) test (see [1, 2, 3] and references therein).

The $K_L \rightarrow \gamma\gamma^*$ form factor (together with $K_L \rightarrow \gamma^*\gamma^*$) is an important ingredient to evaluate properly the dispersive contribution to the real part of the amplitude for the decay $K_L \rightarrow \gamma^*\gamma^* \rightarrow \mu^+\mu^-$. Since this real part receives also short distance contributions proportional to the CKM matrix element V_{td} [4] and the absorptive amplitude is found to saturate the experimental result, a strong cancellation in the real part between short and long distance contributions or the addition of two very small amplitudes is expected.

The importance of $K_L \rightarrow \pi^0\gamma\gamma$ goes further the interest of the process in itself because its role as a CP-conserving two-photon discontinuity amplitude to $K_L \rightarrow \pi^0 e^+e^-$ in possible competition with the CP-violating contributions [2, 3, 5] which are predicted to be of $O(10^{-12})$. The size of the CP-conserving contribution depends on an helicity amplitude for $K_L \rightarrow \pi^0\gamma\gamma$ which appears only at next-to-leading order and this contribution can be controlled both theoretical and experimentally, as we shall see.

The interplay between experimental results and phenomenology indicates that in both these decays ($K_L \rightarrow \pi^0\gamma\gamma$ and $K_L \rightarrow \gamma\gamma^*$) there is an important vector meson exchange contribution. Actually the experimental determination of the slope in $K_L \rightarrow \gamma\gamma^*$ is only based in the BMS

¹To appear in the Proceedings of the XVI International Workshop on Weak Interactions and Neutrinos (WIN'97) Capri, Italy, June 22-28, 1997.

model [6], thus we propose an alternative analysis less model dependent. We realize also that the common problematic point between $K \rightarrow \pi\gamma\gamma$ and $K_L \rightarrow \gamma\gamma^*$ is the model dependence of the weak $VP\gamma$ vertex. Thus after constructing the most general leading χ PT lagrangian for the weak $VP\gamma$ vertex for the processes under consideration we propose a Factorization Model in the Vector couplings (FMV).

2 $K \rightarrow \pi\gamma\gamma$ and $K_L \rightarrow \gamma\gamma^*$ amplitudes

The general amplitude for $K_L(p) \rightarrow \pi^0\gamma(q_1)\gamma(q_2)$ can be written in terms of two independent Lorentz and gauge invariant amplitudes : $A(z, y)$ and $B(z, y)$, where $y = p \cdot (q_1 - q_2)/m_K^2$ and $z = (q_1 + q_2)^2/m_K^2$. Then the double differential rate is given by

$$\frac{\partial^2\Gamma}{\partial y \partial z} = \frac{m_K}{2^9\pi^3} [z^2 |A + B|^2 + \left(y^2 - \frac{\lambda(1, r_\pi^2, z)}{4}\right)^2 |B|^2], \quad (1)$$

where $\lambda(a, b, c)$ is the usual kinematical function and $r_\pi = m_\pi/m_K$. Thus in the region of small z (collinear photons) the B amplitude is dominant and can be determined separately from the A amplitude. This feature is important in order to evaluate the CP conserving contribution $K_L \rightarrow \pi^0\gamma\gamma \rightarrow \pi^0 e^+ e^-$. Both on-shell and off-shell two-photon intermediate states generate, through the A amplitude, a contribution to $K_L \rightarrow \pi^0 e^+ e^-$ that is helicity suppressed [7]. Instead the B -type amplitude, though appearing only at $\mathcal{O}(p^6)$, generates a relevant unsuppressed contribution to $K_L \rightarrow \pi^0 e^+ e^-$ through the on-shell photons [5], due to the different helicity structure.

If CP is conserved the decay $K_L \rightarrow \gamma(q_1, \epsilon_1)\gamma^*(q_2, \epsilon_2)$ is given by an only amplitude $A_{\gamma\gamma^*}(q_2^2)$ that can be expressed as $A_{\gamma\gamma^*}(q_2^2) = A_{\gamma\gamma}^{exp} f(x)$, where $A_{\gamma\gamma}^{exp}$ is the experimental amplitude $A(K_L \rightarrow \gamma\gamma)$ and $x = q_2^2/m_K^2$. The form factor $f(x)$ is properly normalized to $f(0) = 1$ and the slope b of $f(x)$ is defined as $f(x) = 1 + bx$. Traditionally experiments [8] do not measure directly the slope but they input the full form factor suggested by the BMS model [6], however we think that is more appropriate to measure directly the slope and we estimate [9], $b_{exp} = 0.81 \pm 0.18$. The slope b gets two different contributions: the first one (b_V) comes from the strong vector interchange with the weak transition in the K_L leg, the second comes from a direct weak transition $K_L \rightarrow V\gamma$ (b_D). Then $b = b_V + b_D$. While the first term is model independent, the direct contribution b_D requires a modelization due to our ignorance of the weak couplings involving vector mesons. BMS model [6] suggests that the structure of the weak $VP\gamma$ vertex is dominated by a weak vector-vector transition.

The leading finite $\mathcal{O}(p^4)$ amplitudes of $K_L \rightarrow \pi^0\gamma\gamma$ were evaluated some time ago [10], generating only the A -type amplitude in Eq. (1). The observed branching ratio for $K_L \rightarrow \pi^0\gamma\gamma$ is $(1.7 \pm 0.3) \times 10^{-6}$ [8] which is about 3 times larger than the $\mathcal{O}(p^4)$ prediction [10, 11]. However the $\mathcal{O}(p^4)$ spectrum of the diphoton invariant mass nearly agrees with the experiment, in particular no events for small $m_{\gamma\gamma}$ are observed, implying a small B -type amplitude. Thus $\mathcal{O}(p^6)$ corrections have to be important. Though no complete calculation is available, the supposedly larger contributions have been performed : $\mathcal{O}(p^6)$ unitarity corrections [11, 12, 13] enhance the $\mathcal{O}(p^4)$ branching ratio by 30%, and generate a B -type amplitude.

One can parameterize the $\mathcal{O}(p^6)$ vector meson exchange contributions to $K_L \rightarrow \pi^0\gamma\gamma$, by an

effective vector coupling a_V [14] :

$$\begin{aligned} A &= \frac{G_8 M_K^2 \alpha_{em}}{\pi} a_V (3 - z + r_\pi^2) , \\ B &= -\frac{2G_8 M_K^2 \alpha_{em}}{\pi} a_V , \end{aligned} \quad (2)$$

where G_8 is the effective coupling of the leading octet weak chiral lagrangian and is fixed by $K \rightarrow \pi\pi$. Analogously to the $K_L \rightarrow \gamma\gamma^*$ case there are two sources for a_V : i) strong vector resonance exchange with an external weak transition (a_V^{ext}), and ii) direct vector resonance exchange between a weak and a strong $VP\gamma$ vertices (a_V^{dir}). Then $a_V = a_V^{ext} + a_V^{dir}$. The first one is model independent and gives $a_V^{ext} \simeq 0.32$ [14], while the direct contribution depends strongly on the model for the weak $VP\gamma$ vertex. Cohen , Ecker and Pich noticed [12] that one could, simultaneously, obtain the experimental spectrum and width of $K_L \rightarrow \pi^0\gamma\gamma$ with $a_V \simeq -0.9$. The comparison of this result with the value of a_V^{ext} shows the relevance of the direct contributions.

The question of the relevant $\mathcal{O}(p^6)$ contributions relative to the leading $\mathcal{O}(p^4)$ result for $K^+ \rightarrow \pi^+\gamma\gamma$ [7] can also be studied. $\mathcal{O}(p^6)$ unitarity corrections [15] generate a B -type amplitude and increase the rate by a 30–40% while, differently from $K_L \rightarrow \pi^0\gamma\gamma$, vector meson exchange is negligible in $K^+ \rightarrow \pi^+\gamma\gamma$ [9]. BNL-787 has got, for the first time events in this channel [16] confirming the relevance of the unitarity corrections.

3 Factorization Model in the Vector Couplings (FMV)

The general effective weak coupling $VP\gamma$ contributing to $\mathcal{O}(p^6)$ $K \rightarrow \pi\gamma\gamma$ and $K_L \rightarrow \gamma\gamma^*$ processes is

$$\mathcal{L}_W(VP\gamma) = G_8 F_\pi^2 \varepsilon_{\mu\nu\alpha\beta} \sum_{i=1}^5 \kappa_i \langle V^\mu T_i^{\nu\alpha\beta} \rangle , \quad (3)$$

where $T_i^{\nu\alpha\beta}$ $i = 1, \dots, 5$ are all the possible relevant $P\gamma$ structures [9], the κ_i are the dimensionless coupling constants to be determined from phenomenology or theoretical models, $F_\pi \simeq 93$ MeV is the pion decay constant and the brackets in Eq. (3) stand for a trace in the flavour space.

Motivated by the $1/N_c$ expansion the Factorization Model (FM) [17] assumes that the dominant contribution to the four-quark operators of the $\Delta S = 1$ Hamiltonian comes from a factorization *current* $\times$ *current* of them. This assumption is implemented with a bosonization of the left-handed quark currents in χ PT as given by

$$\overline{q_j L} \gamma^\mu q_{iL} \longleftrightarrow \frac{\delta S[U, \ell, r, s, p]}{\delta \ell_{\mu, ji}} , \quad (4)$$

where i, j are flavour indices, and S is the low-energy strong effective action of QCD in terms of the Goldstone bosons realization U and the external fields ℓ, r, s, p .

The general form of the lagrangian is

$$\mathcal{L}_{FM} = 4 k_F G_8 \langle \lambda \frac{\delta S}{\delta \ell_\mu} \frac{\delta S}{\delta \ell^\mu} \rangle + h.c. , \quad (5)$$

where $\lambda \equiv \frac{1}{2}(\lambda_6 - i\lambda_7)$. The FM gives a full prediction but for a fudge factor $k_F \sim \mathcal{O}(1)$.

The procedure used in the FM to study the resonance exchange contribution to a specified process involves first the construction of a resonance exchange generated strong Lagrangian, in terms of Goldstone bosons and external fields, from which to evaluate the left-handed currents. For example, in $K \rightarrow \pi\gamma\gamma$ one starts from the strong $VP\gamma$ vertex and integrates out the vector mesons between two of those vertices giving a strong Lagrangian for $PP\gamma\gamma$ (P is short for pseudoscalar meson) from which to evaluate the left-handed current. This method of implementing the FM imposes as a constraint the strong effective $PP\gamma\gamma$ vertex and therefore it can overlook parts of the chiral structure of the weak vertex. In the FM model the dynamics is hidden in the fudge factor k_F . Alternatively one could try to identify the different vector meson exchange contributions and then estimate the relative weak couplings.

We propose precisely this scheme. Instead of getting the strong lagrangian generated by vector meson exchange we apply the factorization procedure to the construction of the weak $VP\gamma$ vertex and we integrate out the vector mesons afterwards. The interesting advantage of our approach is that, as we have seen in the processes we are interested in, it allows us to identify new contributions to the left-handed currents and therefore to the chiral structure of the weak amplitudes. This we call the Factorization Model in the Vector couplings (FMV).

Hence we obtain the κ_i in Eq. (3) in this framework and we can give a prediction from the FMV model to both a_V^{dir} in $K_L \rightarrow \pi^0\gamma\gamma$ and the slope b_D of $K_L \rightarrow \gamma\gamma^*$ [9]. We get $a_V^{dir}|_{FMV} \simeq -0.95$ and $b_D^{ctet}|_{FMV} \simeq 0.51$. Our final predictions (see Ref. [9] for a thorough discussion) are :

$$a_V \simeq -0.72 \quad , \quad b \simeq 0.8 - 0.9 \quad , \quad (6)$$

in good agreement with phenomenology. Thus we can predict the CP conserving contribution: $0.1 < B(K_L \rightarrow \pi^0 e^+ e^-) \cdot 10^{12} < 3.6$ for $-1.0 < a_V < -0.4$ respectively [9].

4 Dispersive two-photon $B(K_L \rightarrow \mu^+ \mu^-)$

To fully exploit the potential of $K_L \rightarrow \mu^+ \mu^-$ in probing short-distance dynamics it is necessary to have a reliable control on its long-distance amplitude. However the dispersive contribution generated by the two-photon intermediate state cannot be calculated in a model independent way and it is subject to various uncertainties [18]. The branching ratio can be generally decomposed as $B(K_L \rightarrow \mu^+ \mu^-) = |\Re \mathcal{A}|^2 + |\Im \mathcal{A}|^2$, and the dispersive contribution can be rewritten as $\Re \mathcal{A} = \Re \mathcal{A}_{long} + \Re \mathcal{A}_{short}$. The recent measurement of $B(K_L \rightarrow \mu^+ \mu^-)$ [19] is almost saturated by the absorptive amplitude leaving a very small room for the dispersive contribution : $|\Re \mathcal{A}_{exp}|^2 = (-1.0 \pm 3.7) \times 10^{-10}$ or $|\Re \mathcal{A}_{exp}|^2 < 5.6 \times 10^{-10}$ at 90% C.L.

Within the Standard Model the NLO short-distance amplitude [20] gives the possibility to extract a lower bound on $\bar{\rho}$ ($\bar{\rho} = \rho(1 - \lambda/2)$ and ρ and λ the usual Wolfenstein parameters) as

$$\bar{\rho} > 1.2 - \max \left\{ \frac{|\Re \mathcal{A}_{exp}| + |\Re \mathcal{A}_{long}|}{3 \times 10^{-5}} \left[\frac{\overline{m}_t(m_t)}{170 \text{ GeV}} \right]^{-1.55} \left[\frac{|V_{cb}|}{0.040} \right]^{-2} \right\} . \quad (7)$$

In order to saturate this lower bound we propose [21] a low energy parameterization of the $K_L \rightarrow \gamma^* \gamma^*$ form factor that include the poles of the lowest vector meson resonances with arbitrary residues

$$f(q_1^2, q_2^2) = 1 + \alpha \left(\frac{q_1^2}{q_1^2 - m_V^2} + \frac{q_2^2}{q_2^2 - m_V^2} \right) + \beta \frac{q_1^2 q_2^2}{(q_1^2 - m_V^2)(q_2^2 - m_V^2)} . \quad (8)$$

The parameters α and β , expected to be $\mathcal{O}(1)$ by naive dimensional chiral power counting, are in principle directly accessible by experiment in $K_L \rightarrow \gamma \ell^+ \ell^-$ and $K_L \rightarrow e^+ e^- \mu^+ \mu^-$. Up to now there is no information on β and we have the phenomenological result $\alpha = -1.63 \pm 0.22$. The form factor defined in Eq. (8) goes as $1 + 2\alpha + \beta$ for $q_i^2 \gg m_V^2$ and one has to introduce an ultraviolet cutoff Λ . However, in this region the perturbative QCD calculation of the two-photon contribution gives a small result. In particular we find $|1 + 2\alpha + \beta| \ln(\Lambda/M_V) < 0.4$, showing a mild behaviour of the form factor at large q^2 . The FMV model described in the previous Section gives $(1 + 2\alpha + \beta)_{FMV} \simeq -0.01$ in good agreement with the perturbative result.

Using α_{exp} and the QCD constraint we predict [21]

$$|\Re \mathcal{A}_{long}| < 2.9 \times 10^{-5} \quad (90\% C.L.) \quad (9)$$

and

$$\bar{\rho} > -0.38, \quad \text{or} \quad \rho > -0.42 \quad (90\% C.L.) . \quad (10)$$

These bounds could be very much improved if the α and β parameters were measured with good precision and a more stringent bound on $|\Re \mathcal{A}_{exp}|$ is established.

5 Acknowledgement

I wish to thank G. Isidori and J. Portolés for the very fruitful collaborations and discussions.

References

- [1] A. Pich, this Proceedings and refs. therein.
- [2] G. D'Ambrosio, G. Ecker, G. Isidori and H. Neufeld, "Radiative non-leptonic kaon decays" in the Second DAΦNE Physics Handbook, ed. by L. Maiani, G. Pancheri, N. Paver, LNF (1995), p. 265.
- [3] G. D'Ambrosio and G. Isidori, "CP violation in Kaon decays", Preprint INFNNA-IV-96/29, (1996), hep-ph/9611284.
- [4] V.A. Novikov, M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Phys. Rev.*, **D16** (1977) 223.
- [5] J.F. Donoghue and F. Gabbiani, *Phys. Rev.*, **D51** (1995) 2187.
- [6] L. Bergström, E. Massó and P. Singer, *Phys. Lett.*, **B131** (1983) 229, **B249** (1990) 141.
- [7] G. Ecker, A. Pich and E. de Rafael, *Nucl. Phys.*, **B303** (1988) 665.
- [8] Review of Particle Properties, R.M. Barnett et al., *Phys. Rev.*, **D54** (1996) 1.
- [9] G. D'Ambrosio and J. Portolés, *Nucl. Phys.*, **B492** (1997) 417.
- [10] G. Ecker, A. Pich and E. de Rafael, *Phys. Lett.*, **B189** (1987) 363; L. Cappiello and G. D'Ambrosio, *Nuovo Cimento*, **99A** (1988) 155.
- [11] L. Cappiello, G. D'Ambrosio and M. Miragliuolo, *Phys. Lett.*, **B298** (1993) 423.

- [12] A. G. Cohen, G. Ecker and A. Pich, *Phys. Lett.*, **B304** (1993) 347.
- [13] J. Kambor and B.R. Holstein, *Phys. Rev.*, **D49** (1994) 2346.
- [14] G. Ecker, A. Pich and E. de Rafael, *Phys. Lett.*, **B237** (1990) 481.
- [15] G. D'Ambrosio and J. Portolés, *Phys. Lett.*, **B389** (1996) 770; (E) *ibid.*, **B395** (1997) 389.
- [16] P. Kitching et al., “Observation of the decay $K^+ \rightarrow \pi^+ \gamma \gamma$ ”, Preprint BNL-64628 (1997), hep-ex/9708011.
- [17] A. Pich and E. de Rafael, *Nucl. Phys.*, **B358** (1991) 311.
- [18] M.B. Voloshin and E.P. Shabalin, *JETP Lett.*, **23** (1976) 107; R.E. Shrock and M.B. Voloshin, *Phys. Lett.*, **B87** (1980) 375; L. Bergström, E. Massó and P. Singer, *Phys. Lett.*, **B131** (1983) 229; G. D'Ambrosio and D. Espriu, *Phys. Lett.*, **B175** (1986) 237; G. Belanger and C.Q. Geng, *Phys. Rev.*, **D43** (1991), 140; P. Ko, *Phys. Rev.*, **D45** (1992) 174.
- [19] A.P. Heinson et al., *Phys. Rev.*, **D51** (1995) 985.
- [20] G. Buchalla and A.J. Buras, *Nucl. Phys.*, **B412** (1994) 106.
- [21] G. D'Ambrosio, G. Isidori and J. Portolés, “Can we extract short-distance information from $B(K_L \rightarrow \mu^+ \mu^-)$?”, Preprint INFNNA-IV-97/40 (1997), hep-ph/9708326.